

TABLE 1.

Clinical Characteristics, Treatment, and Response to Treatment in Cases of LPP						
Age at Onset/ Gender	Scalp Involvement (Symptoms and Location)	Non-scalp Involvement (Symptoms and Location)	Therapy	Response to Therapy	Follow-Up Time (Months)	Reference Number
15/M	Scarring alopecia of the parietal and occipital scalp; erythema and perifollicular scaling on the plaques' periphery with erythematous, shiny central areas with notable absence of follicular ostia	None	Clobetasol propionate (0.5 mg/g) gel to affected areas, twice daily for three months	Partial Response	Not reported	3
6/F	Frontal and parietal scalp involvement with perifollicular papules	Lateral eyebrow alopecia Bilateral ocular involvement: extensive corneal scarring, epithelial breakdown and vascularization of right and left eyes that progressed to complete opacification and hypertrophic scarring bilaterally	Cyclosporine (40 mg twice daily)	Not reported	Not reported	4
13/M	Pruritus and perifollicular scale involving the vertex and parietal scalp	Nail Pitting	High potency topical corticosteroid; Intralesional triamcinolone acetonide (40 mg/mL)	Not Reported	Not Reported	5
14/F	Right posterior scalp involvement; Scarring, atrophy, erythema	None	Topical clobetasol propionate (0.05%); Intralesional triamcinolone acetonide (0.3 mL of 40 mg/mL; 10 injections over 1.5 years)	Partial response	17	5
14/M	Superior and lateral scalp involvement characterized by perifollicular erythema and scale	Nail Ridging	Minocycline (100 mg twice daily for one year)	Complete Response	60	5
16/M	Perifollicular erythema and scale limited to the vertex scalp	None	Topical clobetasol propionate (0.05% twice daily)	Complete Response	3	5
16/F	Pruritus and scarring limited to the vertex scalp	None	Topical Corticosteroid	Complete Response	3	6
9/F	Pruritus and scarring limited to the scalp vertex scalp	None	Topical Corticosteroid	Partial Response	6	6
8/M	Frontal scalp alopecia characterized by perifollicular inflammation and follicular plugging	None	Topical Corticosteroid; Topical Tacrolimus	No response	24	7
12/M	Multicentric alopecia with extensive hyperkeratotic follicular papules	Erythema dyschromicum perstans	Griseofulvin (12.5 mg/kg)	No response	3	8
12/F	Vertex and temporal scalp involvement characterized by perifollicular inflammation	Skin involvement	Topical clobetasol propionate (0.05%) followed by sulfasalazine, which was subsequently replaced by tocilizumab to treat oligoarthritis	Complete response	12	9
10/M	Vertex scalp involvement; scarring alopecia with perifollicular erythema and keratin plugs	Skin involvement	Not Available	Not Reported	Not Available	10
12/M	Scarring alopecia limited to the vertex scalp; prominent erythema, scaly follicular papules and shiny, atrophic skin	Skin Involvement	Griseofulvin (375 mg/d)	Complete Response	6	11

F= Female; LPP= Lichen Planopilaris; M= Male